



ELSEVIER

Brazilian Journal of  
OTORHINOLARYNGOLOGY

[www.bjorl.org](http://www.bjorl.org)



ORIGINAL ARTICLE

**Association between interleukin-6 polymorphism in the –174 G/C region and hearing loss in the elderly with a history of occupational noise exposure<sup>☆,☆☆</sup>**

Miula Portelinha Braga<sup>a</sup>, Sandra Mara Maciel<sup>b,c</sup>, Luciana Lozza de Moraes Marchiori<sup>d,a,\*</sup>,  
Regina Célia Poli-Frederico<sup>a,e</sup>

<sup>a</sup> Universidade Norte do Paraná (UNOPAR), Londrina, PR, Brazil

<sup>b</sup> University of London, London, United Kingdom

<sup>c</sup> Department of Pediatric Dentistry, Universidade Estadual de Maringá (UEM), Maringá, PR, Brazil

<sup>d</sup> Universidade Estadual de Londrina (UEL), Londrina, PR, Brazil

<sup>e</sup> Universidade Estadual Paulista "Júlio de Mesquita Filho" (UNESP), Botucatu, PR, Brazil

Received 29 July 2013; accepted 1 February 2014

Available online 22 July 2014

**KEYWORDS**

Occupational noise;  
Sensorineural hearing  
loss;  
Cytokines

**Abstract**

**Introduction:** The biological processes involved in noise-induced hearing loss (NIHL) are still unclear. The involvement of inflammation in this condition has been suggested.

**Objective:** To investigate the association between interleukin – 6 (IL-6) polymorphism and susceptibility to NIHL.

**Methods:** This was a cross-sectional study with a sample of 191 independent elderly individuals aged >60 years of age. Information on exposure to occupational noise was obtained by interviews. Audiological evaluation was performed using pure tone audiometry and genotyped through PCR by restriction fragment length polymorphism – PCR-RFLP. Data were analyzed using the chi-square test and the odds ratio (OR), with the significance level set at 5%.

**Results:** Among elderly with hearing loss (78.0%), 18.8% had a history of exposure to occupational noise. There was a statistically significant association between the genotype frequencies of the IL-6 –174 and NIHL. The elderly with the CC genotype were less likely to have hearing loss due to occupational noise exposure when compared to those carrying the GG genotype (OR = 0.0124; 95% CI 0.0023–0.0671;  $p < 0.001$ ).

**Conclusion:** This study suggests there is an association of polymorphisms in the IL-6 gene at position – G174C with susceptibility to noise-induced hearing loss.

© 2014 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Published by Elsevier Editora Ltda. All rights reserved.

<sup>☆</sup> Please cite this article as: Braga MP, Maciel SM, Marchiori LL, Poli-Frederico RC. Association between interleukin-6 polymorphism in the –174 G/C region and hearing loss in the elderly with a history of occupational noise exposure. Braz J Otorhinolaryngol. 2014;80:373–8.

<sup>☆☆</sup> Institution: Universidade Norte do Paraná (UNOPAR), Londrina, PR, Brazil.

\* Corresponding author at: Universidade Norte do Paraná (UNOPAR), Londrina, PR, Brazil.

E-mail: [luciana.marchiori@unopar.br](mailto:luciana.marchiori@unopar.br) (L.L.M. Marchiori).

## PALAVRAS-CHAVE

Ruído ocupacional;  
Perda auditiva  
neurosensorial;  
Citocinas

## Associação entre interleucina-6 polimorfismo na região de -174 G/C e perda auditiva em idosos com história de exposição ao ruído ocupacional

### Resumo

**Introdução:** Os processos biológicos envolvidos na perda auditiva induzida por ruído (PAIR) ainda não estão claros. O envolvimento de processo inflamatório nesta condição tem sido sugerido.

**Objetivo:** Investigar a associação entre o polimorfismo no gene da interleucina-6 (IL-6) e a suscetibilidade à PAIR.

**Método:** Trata-se de estudo transversal com amostra de 191 idosos independentes acima de 60 anos de idade. Informações sobre a exposição ao ruído ocupacional foram obtidas por entrevistas. A avaliação audiológica foi realizada por meio de audiometria tonal liminar e a genotipagem pela técnica da PCR-RFLP. Os dados foram analisados usando-se o teste Qui-quadrado e a razão de chances (OR), com o nível de significância fixado em 5%.

**Resultados:** Entre os idosos com perda auditiva (78,0%), 18,8% apresentavam histórico de exposição ao ruído ocupacional. Houve associação estatisticamente significante entre as frequências genotípicas da IL-6 -174 e a PAIR. Os idosos portadores do genótipo CC foram menos propensos a apresentar perda auditiva por exposição ao ruído ocupacional quando comparados a aqueles portadores do genótipo GG (OR = 0,0124; 95% IC 0,0023-0,0671; p < 0,001).

**Conclusão:** O presente estudo sugere a associação do polimorfismo no gene da IL-6 na posição -G174C com a suscetibilidade à perda auditiva induzida por ruído.

© 2014 Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

## Introduction

Elderly health has increasingly aroused the interest of researchers, as the population in both developed and developing countries is aging.<sup>1</sup> It is estimated that by 2040 developing countries will have a billion people aged 60 years or older.<sup>2</sup> Given the rapidity and magnitude of this increase, care for this specific group is essential, so they can age in good health and with good quality of life.<sup>3</sup>

The most common types of hearing loss are presbycusis (due to aging) and noise-induced hearing loss (NIHL).<sup>4</sup> NIHL is due to continued exposure to loud noise levels, resulting in gradual loss of auditory acuity and is usually bilateral, symmetric, sensorineural and irreversible. It usually affects the high audiometric frequencies of 3000–6000 Hz.<sup>5</sup> The etiology of hearing loss also has genetic components, in addition to environmental ones, and more recently, inner ear inflammatory response induction and up-regulation of proinflammatory cytokines were evaluated in the presence of exacerbated noise.<sup>5–7</sup> Several researchers have reported the presence of inflammatory cells in the steady state and their increase after lesions in the inner ear.<sup>6,8–10</sup>

Noise exposure induces the expression of pro-inflammatory cytokines, including tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin 1 $\beta$  (IL-1 $\beta$ ) and interleukin-6 (IL-6).<sup>11</sup> So et al.<sup>12</sup> observed a transient up-regulation of IL-6 in cisplatin-treated animal models. Wakabayashi et al.<sup>13</sup> investigated the effect of IL-6 inhibition using an anti-IL-6 (MR16-1) antibody in mice. These authors found that MR16-1 showed a protective effect against noise-induced cochlear injury, mainly due to neuronal loss suppression and presumably by relieving the inflammatory response, similar data were found by Nakamoto et al.<sup>14</sup> These authors suggested that the suppression of proinflammatory

cytokine HSF-1 in the cochlea by the administration of GGA (geranyl-geranyl-acetone) could be an important way to protect the inner ear.

The expression of cytokines may be influenced by genetic variation, resulting in pathogenic conditions<sup>15</sup> and several studies have investigated single nucleotide polymorphisms (SNPs) as risk factors for inflammatory diseases.<sup>16</sup> These SNPs may affect the expression, secretion and cellular transport of interleukins<sup>17,18</sup> and can also decrease the level of the IL-1 receptor antagonist (IL-1Ra), which increases the production and activity of IL-1 $\beta$ .<sup>19</sup>

The SNP in the region –174 of the interleukin-6 gene contains the replacement of G by C, and the presence of the C allele represents a protective function due to reduced production of IL-6.<sup>20</sup>

Considering that NIHL is involved in the inflammatory process, this study aimed to evaluate the association between the proinflammatory cytokine-6 (IL-6) gene polymorphism and susceptibility to NIHL in physically independent elderly Brazilians.

## Method

### Study population

This cross-sectional study was approved by the Ethics Committee on Human Research of the University of Northern Paraná (0070/09). It is part of a broader investigation, the “Study on aging and longevity”, which has been carried out in Londrina since 2009. Londrina (approximately 500,000 inhabitants) is located in the northern region of the state of Paraná, Brazil.

Of a population of 43,610 elderly individuals treated in 38 basic health units (BHUs) in the urban area of the city, the sample size was set at 343 individuals, considering a confidence interval of 95% and a margin of error of 5%.<sup>21</sup>

Aiming at obtaining sample representativeness, random stratification was performed considering gender and regions of the city. The study included individuals aged 60 years or older, of both genders, living independently and classified at level 3 or 4, as proposed by Spirduso.<sup>22</sup> This classification assesses the level of independence of the elderly, in which level 1 indicates a lack of self-mobility and level five indicates athletes. The elderly who had some disease or limitations that prevented the tests, such as physical or mental impairment, were excluded from the sample. All participants signed an informed consent form.

## Audiological assessment

The audiological assessment was performed by conventional pure tone audiometry, in an interacoustics audiometer. To determine the severity of hearing loss, the criteria proposed by BIAP – *Bureau International d'Audio Phonologie* – an institution formed by several associations of European countries with the main objective of guiding the activity of professionals in these regions, were used. The means of hearing thresholds in dB in air conduction, at 500 Hz, 1 kHz, 2 kHz and 4 kHz and the 02/1 1997 recommendations were analyzed, both the right and left ears were evaluated for a group of individuals with a history of occupational noise and the group of individuals with no history of occupational noise.<sup>23-25</sup>

## Evaluation of occupational noise exposure

The assessment of occupational noise exposure was performed through interviews with the elderly participants, using a semi-structured questionnaire. Information was collected on whether work was performed in a noisy environment or not, how many years of work in a noisy environment and whether the individual wore a hearing aid. Moreover, demographic characteristics (gender, age and ethnicity) were collected.

## Genotyping for IL-6 G-174C (rs1800795) gene polymorphism

Peripheral blood samples were collected in vacuum tubes containing 6% ethylenediamine tetraacetic acid (EDTA) and DNA was extracted from peripheral blood leukocytes, using the protocol described by Olerup and Zeterquist.<sup>26</sup>

Based on the original sample, the DNA was diluted to a concentration of 100 ng/mL. The concentration was determined by spectrophotometer at 260 nm and 280 nm (Biomate 3, Thermo Fisher Scientific, Madison, USA).

The C to T polymorphic site located at position -174 of the IL-6 (rs1800795) gene was amplified by polymerase chain reaction (PCR) resulting in a 99 bp fragment. The PCR mixture contained a mixture of a buffer solution pH 8.0 1×, 1.5 mM of MgCl<sub>2</sub>, 8 mM each of deoxyribonucleotide triphosphates, 1 μM of each primer and 1 U Taq DNA

Polymerase (Invitrogen, Carlsbad, CA, USA). The primers used were F: 5'-TTGTCAAGACATGCCAAGTGCT-3' and R: 5'-GCCTCAGAGACATCTCCAGTCC-3' (Invitrogen, Carlsbad, CA, USA).<sup>27</sup>

PCR amplification was performed in a thermocycler (TC-512 Techne resistance, Burlington, NJ, USA) under the following conditions: initial denaturation at 95 °C for 5 min followed by 30 cycles of 95 °C for 1 min, 56 °C for 1 min and 72 °C for 1 min, and final extension at 72 °C for 5 min.

The PCR product was digested with 2 U of restriction enzyme NtallI (Invitrogen, Carlsbad, CA, USA) overnight at 65 °C. The digested fragments were separated on 2% agarose gel (Invitrogen Life Technologies, São Paulo, Brazil). A sample with known genotype was used as positive control and ultra-pure water was used as negative PCR control.

The 100-bp DNA molecular weight marker (Ladder, Invitrogen) was included on each gel stained with SybrSafe (Invitrogen Life Technologies, São Paulo, Brazil) and visualized under UV illumination. The reading and interpretation of agarose gels were made with LabImage L-PIX (HE) 1D-L340 (Loccus Biotechnology, São Paulo, Brazil) program. Fragments of 13, 227 and 59 bp (allele G), and fragments of 13, 118, 109 and 59 bp (allele C) were observed.

## Statistical analysis

The statistical analysis was performed using the statistical package SPSS 17.0 (SPSS Inc., Chicago, IL, USA). To test the association between genotype frequency and hearing loss related to a history of occupational noise exposure, the chi-square test was performed. The Hardy-Weinberg equilibrium was tested in each group using the chi-square test. The odds ratio (OR) with a confidence interval of 95% (95% CI) was calculated. The significance level was set at 5%.

## Results

Of a total of 343, the molecular genetic procedures were performed in 191 elderly. Of these, 31.9% reported a history of noise exposure, with 18.8% exhibiting hearing impairment. Of the 68.1% with no history of noise exposure, 59.2% had hearing loss. The mean age was  $67.8 \pm 5.3$  years, with a higher proportion (58.6%) of elderly women (Table 1).

Of the assessed elderly individuals, 56% were homozygous for the G allele, 7.8% were homozygous for the C allele and 36.2% were heterozygous (Table 1). The genotype distribution for IL-6 gene was in agreement with the Hardy-Weinberg equilibrium ( $p > 0.05$ ).

A statistically significant association was found between the genotype and allele frequencies of the IL-6 -174 (rs1800795) gene and hearing loss among elderly adults with a history of occupational noise exposure ( $p < 0.001$ ) (Table 2). The elderly with the CC genotype were less likely to have hearing loss due to occupational noise exposure in relation to the elderly with the GG genotype (OR = 0.0124; 95% CI: 0.0023–0.0671;  $p < 0.0001$ ). However, this association was not observed in the elderly group without a history of noise exposure ( $\chi^2 = 0.078$ ;  $p = 0.961$ ).

**Table 1** General characteristics, allelic and genotypic frequencies among Brazilian elderly ( $n=191$ ).

Characteristics	n	%
<i>Gender</i>		
Male	79	41.4
Female	112	58.6
<i>Age (years)</i>		
60–64	63	33.0
65–74	104	54.5
75 or +	24	12.6
<i>Mean 67.8 (SD = 5.3)</i>		
<i>History of noise exposure</i>		
No history of noise exposure		
No hearing loss	17	8.9
With hearing loss	113	59.2
<i>With history of noise exposure</i>		
No hearing loss	25	13.1
With hearing loss	36	18.8
<i>Genotypic frequency</i>		
GG	107	56.0
CC	15	7.8
GC	69	36.2
<i>Allelic frequency</i>		
G	283	74.1
C	99	25.9

## Discussion

Although the mechanism and role of proinflammatory cytokines in NIHL are not well understood, it is known

that the structure and expression of cytokines may be affected by genetic variation, such as single nucleotide polymorphisms (SNPs), which results in evident pathological consequences.<sup>15</sup> SNP functionality regarding gene expression is an important subject in disease-associated studies.<sup>16</sup>

In the present study, the frequencies of alleles G and C in the study population were 74.1% and 25.9%, respectively. Other studies have also demonstrated a predominance of the ancestral allele (G) ranging from 60% to 70%.<sup>28</sup> Furthermore, we observed a higher frequency of GG genotype (56%), followed by GC (36.2%) and CC (7.8%). This is consistent with the aforementioned studies.<sup>20,28</sup>

We observed an association between the GG genotype of IL-6 and hearing loss with history of exposure to occupational noise ( $\chi^2 = 40.201$ ;  $p < 0.001$ ). The elderly with the CC genotype had lower susceptibility to hearing loss due to occupational noise exposure (OR = 0.0124; 95% CI: 0.0023–0.0671;  $p < 0.01$ ) compared to elderly patients with the GG genotype. The presence of the C allele results in lower expression of IL-6 after an inflammatory stimulus, when compared with the G allele.<sup>29</sup> It is suggested that the CC genotype confers a protective effect against the development of comorbidities, which can be verified in this analysis, in which the frequency the C allele in the hearing loss group without exposure to noise was higher than in the NIHL group.

The studies by Hirose et al.<sup>6</sup> and Fujioka et al.<sup>11</sup> pointed out the possibility of inflammatory changes in the cochlea after noise stimuli. Fujioka et al.<sup>11</sup> demonstrated for the first time the induction of pro-inflammatory cytokines in the cochlea exposed to the noise and observed increased expression of interleukins TNF- $\alpha$ , IL-1 $\beta$  and IL-6 3 h after noise exposure. These authors pointed out that this is a self-protection mechanism against exposure to large amounts of noise and that the consequent over-expression of interleukins for long periods must worsen cochlear function.

**Table 2** Association between genotype frequency for the interleukin-6 G-174C gene polymorphism and hearing loss related to history of occupational noise exposure ( $n=191$ ).

Occupational noise exposure	Genotyping	No hearing loss n (%)	With hearing loss n (%)	p value
No history	<i>Genotypic frequency</i>			0.961 <sup>a</sup>
	GG	09 (52.9)	62 (54.9)	
	CC	01 (05.9)	08 (07.0)	
	GC	07 (41.2)	43 (38.1)	
	<i>Allelic frequency</i>			0.869 <sup>b</sup>
	G	25 (73.5)	167 (76.9)	
	C	09 (26.5)	59 (23.1)	
With history	<i>Genotypic frequency</i>			<0.01 <sup>c</sup>
	GG	03 (12.0)	33 (91.7)	
	CC	04 (16.0)	02 (05.5)	
	GC	18 (72.0)	01 (02.8)	
	<i>Allelic frequency</i>			<0.01 <sup>d</sup>
	G	24 (48.0)	67 (93.0)	
	C	26 (52.0)	05 (07.0)	

<sup>a</sup>  $\chi^2 = 0.078$ .

<sup>b</sup>  $\chi^2 = 0.027$ .

<sup>c</sup>  $\chi^2 = 40.201$ .

<sup>d</sup>  $\chi^2 = 31.605$ .

The findings by So et al.<sup>12</sup> clearly demonstrated that IL-6 acts as an inducer of acute cochlear damage, considering that a transient up-regulation of IL-6 occurred after exposure of animals with cisplatin. Some studies have shown that the inhibition of IL-6 production has a cochlear protective effect. Wakabayashi et al.<sup>21</sup> used anti-IL-6 (MR10-1) antibody to inhibit IL-6 production in mice, identifying a protective effect for noise-induced cochlear injury. Along this same line of investigation, Nakamoto et al.<sup>14</sup> found that the drug geranyl-geranyl-acetone (GGA) stimulates expression of the HSF gene which, in turn, inhibits inflammation in the cochlea, protecting the inner ear.

It is noteworthy that this strong evidence of the involvement of IL-6 with NIHL has been verified by several studies in animal models.<sup>11-14</sup> Evidence in human beings related to pro-inflammatory interleukins and NIHL was recently reported by our group, when we investigated the association of polymorphisms in the IL-1 $\beta$  gene and NIHL.<sup>30</sup> That study demonstrated that the polymorphism in this gene is not associated with NIHL in the assessed elderly subjects. However, the findings of the present study showed that polymorphisms in the IL-6 gene should contribute to the risk of NIHL in the Brazilian elderly.

Considering this context, the identification of the polymorphism in the IL-6 gene in patients with a history of hearing loss related to occupational exposure to noise may help us to understand individual variability of inflammation resulting in hearing loss, as well as, in the future, to suggest genotyping individuals for this particular polymorphism (rs1800795) in order to determine individual susceptibility, providing a new strategy for the prevention of hearing loss related to noise exposure.

## Conclusions

An association between the presence of IL-6 gene polymorphism and hearing loss associated with occupational noise exposure was found in elderly Brazilians. Studies based on large populations and with different ethnicities should be performed to confirm our findings.

## Funding

This study was funded by FUNADESP / Brazil (Fundação Nacional de Desenvolvimento do Ensino Superior Particular).

## Conflicts of interest

The authors declare no conflicts of interest.

## References

1. Shrestha LB. Population aging in developing countries. *Health Affairs*. 2000;19:204-12.
2. Kinsella K, He W. An aging world: 2008. USA: Census Bureau, Washington; 2009.
3. Veras R. Population aging today: demands, challenges and innovations. *Rev Saude Pública*. 2009;43:548-54.
4. Nudelmann AA, Costa EA, Seligman J, Ibañez RN. Perda auditiva induzida pelo ruído. Porto Alegre: Bagagem; 1997.
5. Keithley EM, Wang X, Barkdull GC. Tumor necrosis factor alpha can induce recruitment of inflammatory cells to the cochlea. *Otol Neurotol*. 2008;29:854-9.
6. Hirose K, Discolo CM, Keasler JR, Ransohoff R. Mononuclear phagocytes migrate into the murine cochlea after acoustic trauma. *J Comp Neurol*. 2005;489:180-94.
7. Satoh H, Firestein GS, Billings PB, Harris JP, Keithley EM. Tumor necrosis factor-alpha, an initiator, and etanercept, an inhibitor of cochlear inflammation. *Laryngoscope*. 2002;112:1627-34.
8. Jokay I, Papp Z, Soos G, Sziklai I, Dezso B. The effect of chronic otitis media on the immunoreactivity of human inner ear. *Eur Arch Otorhinolaryngol*. 2001;258:529-32.
9. Okano T, Nakagawa T, Kita T, Kada S, Yoshimoto M, Nakahata T, et al. Bone marrow-derived cells expressing Iba1 are constitutively present as resident tissue macrophages in the mouse cochlea. *J Neurosci Res*. 2008;86:1758-67.
10. Tan BT, Lee MM, Ruan R. Bone marrow-derived cells that home to acoustic deafened cochlea preserved their hematopoietic identity. *J Comp Neurol*. 2008;509:167-79.
11. Fujioka M, Kanzaki S, Okano HJ, Masuda M, Ogawa K, Okano H. Proinflammatory cytokines expression in noise-induced damaged cochlea. *J Neurosci Res*. 2006;83:575-83.
12. So H, Kim H, Lee JH, Park C, Kim Y, Kim E, et al. Cisplatin cytotoxicity of auditory cells requires secretions of proinflammatory cytokines via activation of ERK and NFκB. *J Assoc Res*. 2007;8:338-55.
13. Wakabayashi K, Fujioka M, Kanzaki S, Okano HJ, Shibata S, Yamashita D, et al. Blockade of interleukin-6 signaling suppressed cochlear inflammatory response and improved hearing impairment in noise-damaged mice cochlea. *Neurosci Res*. 2010;66:345-52.
14. Nakamoto T, Mikuriya T, Sugahara K, Hirose Y, Hashimoto T, Shimogori H, et al. Geranylgeranylacetone suppresses noise-induced expression of proinflammatory cytokines in the cochlea. *Auris Nasus Larynx*. 2012;39:270-4.
15. Satoh H, Firestein GS, Billings PB, Harris JP, Keithley EM. Proinflammatory cytokine expression in the endolymphatic sac during inner ear inflammation. *J Assoc Res Otolaryngol*. 2003;4:139-47.
16. Dinarello CA. Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood*. 2011;117:3720-32.
17. Hwang JH, Chen JC, Yang SY, Wang MF, Chan YC. Expression of tumor necrosis factor- $\alpha$  and interleukin-1 $\beta$  genes in the cochlea and inferior colliculus in salicylate-induced tinnitus. *J Neuroinflamm*. 2011;8:30.
18. Santtila S, Savinainen K, Hurme M. Presence of the IL-1RA allele 2 (IL1RN\*2) is associated with enhanced IL-1 $\beta$  production *in vitro*. *Scand J Immunol*. 1998;47:195-8.
19. Bioque G, Crusius JB, Koutroubakis I, Bouma G, Kostense PJ, Meuwissen SG, et al. Allelic polymorphism in IL-1 beta and IL-1 receptor antagonist (IL-1Ra) genes in inflammatory bowel disease. *Clin Exp Immunol*. 1995;102:379-83.
20. Costa AM, Guimarães MCM, De Souza ER, Nóbrega OT, Bezerra ACB. Interleukin-6 (G-174C) and tumour necrosis factor-alpha (G-308A) gene polymorphisms in geriatric patients with chronic periodontitis. *Gerodontology*. 2010;27:70-5.
21. Raosoft Sample Size Calculator [Internet]. Available from: <http://www.raosoft.com/samplesize.html> (accessed 10.06.13).
22. Spirduso WW. Physical dimensions of aging. 2nd ed. Barueri: Manole; 2005. p. 75-80.
23. National Committee on Noise and Hearing Conservation. Recommendations for the assessment of damage caused by Noise Induced Hearing Loss. *Acta AWHO*. 1996;16:45 [Letter to the Editors].
24. Amorim RB, Lopes AC, Santos KT, Melo AD, Lauris JR. Hearing alterations in the occupational exposure in musicians. *Arch Otorrinolaryngol Int*. 2008;12:377-83.

25. Manual de Procedimentos em Audiometria Tonal Limiar, Logoaudiometria e Medidas de Imitância Acústica. <http://www.fonoaudiologia.org.br/publicacoes/ManualdeAudiologia.pdf> (accessed 18.04.13).
26. Olerup O, Zetterquist H. HLA-DR typing by PCR amplification with sequence-specific primers (PCR-SSP) in 2 hours: an alternative to serological DR typing in clinical practice including donor-recipient matching in cadaveric transplantation. *Tissue Antigens*. 1992;39:225-35.
27. Olomolaiye O, Wood NA, Bidwell JL. A novel NlalII polymorphism in the human IL-6 promoter. *Eur J Immunogenet*. 1998;25:267.
28. Trevilatto PC, Scarel-Caminaga RM, De Brito Jr RB, De Souza AP, Line SRP. Polymorphism at the position -174 of IL-6 is associated with susceptibility to chronic periodontitis in a Caucasian Brazilian population. *J Clin Periodontol*. 2003;30: 438-42.
29. Fishman D, Faulds G, Jeffey R, Mohamed-Ali Y, Yudkin JS, Humphries S, et al. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *J Clin Investig*. 1998;102:1369-76.
30. Carvalho LCL, Marchiori LM, Melo JJ, Maciel SM, Poli-Frederico RC. Interleukin-1 $\beta$  gene polymorphism and hearing loss related to the history of occupational noise exposure in Brazilian elderly. *Noise Health*. 2013;15:160-4.